

**ELIGIBLE PATIENTS WITH COMMERCIAL
INSURANCE MAY PAY AS LITTLE AS**

\$0*

 **THYQUIDITY™**
(levothyroxine sodium) oral solution, 100 mcg/5 mL

ELIGIBLE PATIENTS WITH COMMERCIAL
INSURANCE MAY PAY AS LITTLE AS

\$0*

 **THYQUIDITY™**
(levothyroxine sodium) oral solution, 100 mcg/5 mL

BIN# 610524
PCN# LOYALTY
GROUP# 50777459
ID# 1396558902

SAVINGS CARD

Please visit www.Thyquidity.com
for Indication and Important Safety
Information including Boxed Warning.
Please see full Prescribing Information.

*Must meet eligibility requirements. Azurity
Pharmaceuticals, Inc. reserves the right to
rescind, revoke, or amend this offer without
notice. See Terms and Conditions below.

***Eligible patients with commercial insurance only (non-government),
may use this card at their pharmacy to save on THYQUIDITY prescriptions.**

**See full terms and conditions also available at
www.thyquidity.com/patientresources.**

Please see Important Safety Information Including Boxed Warning on page 3.

Please see full Prescribing information.



THYQUIDITY™

(levothyroxine sodium) oral solution, 100 mcg/5 mL

*Copay Savings Program Terms and Conditions

To the Patient:

If your copay for Thyquidity exceeds \$0 (insured patients), present this card to the pharmacist for an instant rebate. Benefit limitations apply. Patient is responsible for the remaining balance after benefit limits are reached. For questions regarding your eligibility or benefits or if you wish to discontinue your participation, call 1-877-264-2440 (8:00 AM-8:00 PM ET, Monday-Friday).

To the Pharmacist:

Benefit limitations apply. Additional program details are available at www.thyquidity.com. When you use this card, you are certifying that you have not submitted and will not submit a claim for reimbursement under and federal, state, or other governmental programs for this prescription. By redeeming this coupon, you agree that you understand and will abide by the terms and conditions of this offer, posted at www.mckesson.com/mprstnc.

- **Submit transaction to McKesson Corporation using BIN #610524**
- **Patient not eligible if prescriptions are paid in part or full by any state or federally funded programs, including but not limited to, Medicare or Medicaid, Medigap, VA, DOD, or Tricare. This program is not valid where prohibited by law.**
- If primary coverage exists, input card information as secondary coverage and transmit using the COB segment of the NCPDP transaction. Applicable discounts will be displayed in the transaction response.
- Acceptance of this card and your submission of claims for the Thyquidity Savings Card program are subject to the Thyquidity Savings Card Program Terms and Conditions posted at www.mckesson.com/mprstnc.
- LoyaltyScript is not an insurance card.
- For questions regarding setup, claim transmission, patient eligibility or other issues, call 1-877-264-2440 (8:00 AM-8:00PM ET, Monday-Friday).

**Please see Indication and Important Safety Information including Boxed Warning.
Please see full Prescribing Information.**

Not intended for HCPs in Vermont



Hypothyroidism

As a replacement therapy in primary (thyroidal), secondary (pituitary), and tertiary (hypothalamic) congenital or acquired hypothyroidism.

Pituitary Thyrotropin (Thyroid Stimulating Hormone, TSH) Suppression

As an adjunct to surgery and radioiodine therapy in the management of thyrotropin-dependent well-differentiated thyroid cancer.

Limitations of Use

- Not indicated for suppression of benign thyroid nodules and nontoxic diffuse goiter in iodine-sufficient patients.
- Not indicated for treatment of hypothyroidism during the recovery phase of subacute thyroiditis.

IMPORTANT SAFETY INFORMATION**WARNING: NOT FOR TREATMENT OF OBESITY OR FOR WEIGHT LOSS**

See full prescribing information for complete boxed warning

- **Thyroid hormones, including THYQUIDITY, should not be used for the treatment of obesity or for weight loss.**
- **Doses beyond the range of daily hormonal requirements may produce serious or even life-threatening manifestations of toxicity. (6, 10).**

Do not use THYQUIDITY if you have uncorrected adrenal problems.

Thyroid hormones should be used with great caution in circumstances where the integrity of the cardiovascular system is suspected. In the elderly and in patients with cardiovascular disease, THYQUIDITY should be used with greater caution.

Tell your doctor about any other medical conditions you may have, especially heart disease, diabetes, blood clotting problems, and adrenal or pituitary gland problems. The dose of other drugs you may be taking to control these conditions may have to be changed while you are taking THYQUIDITY. If you have diabetes, check your blood sugar levels and/or the glucose in your urine, as ordered by your doctor, and immediately tell your doctor if there are any changes. If you are taking blood thinners, your blood clotting status should be checked often.

Taking too much levothyroxine has been associated with increased bone loss, especially in women after menopause.

Once your doctor has found your specific THYQUIDITY dose, it is important to have lab tests done, as ordered by your doctor, at least once a year.

Foods like soybean flour, cottonseed meal, walnuts, and dietary fiber may cause your body to absorb less THYQUIDITY from the gastrointestinal tract. Grapefruit juice may cause your body to absorb less levothyroxine and may reduce its effect. Let your doctor know if you eat these foods, as your dose of THYQUIDITY may need to be adjusted.

Use THYQUIDITY only as ordered by your doctor. Take THYQUIDITY as a single dose, preferably on an empty stomach, one-half to one hour before breakfast.

Products such as iron and calcium supplements and antacids can lower your body's ability to absorb levothyroxine, so THYQUIDITY should be taken 4 hours before or after taking these products.

Tell your doctor if you are pregnant or breastfeeding or are thinking of becoming pregnant while taking THYQUIDITY. Your dose of THYQUIDITY may need to be increased during your pregnancy.

Monitor your baby from birth to 3 months of age for vomiting and/or diarrhea as THYQUIDITY can cause gastrointestinal irritation due to the glycerol component.

It may take several weeks before you notice an improvement in your symptoms.

Tell your doctor about all medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements.

Tell your doctor or dentist that you are taking THYQUIDITY before any surgery.

Adverse reactions associated with THYQUIDITY therapy are primarily those of hyperthyroidism due to therapeutic overdosage: rapid or abnormal heartbeat, chest pain, difficulty catching your breath, leg cramps, headache, nervousness, irritability, sleeplessness, shaking, change in appetite, weight gain or loss, vomiting, diarrhea, increased sweating, difficulty tolerating heat, fever, changes in menstrual periods, swollen red bumps on the skin (hives) or skin rash, or any other unusual medical event. Tell your doctor if you develop any of these symptoms.

Partial hair loss may occur during the first few months you are taking THYQUIDITY.

The Important Safety Information does not include all the information needed to use THYQUIDITY safely and effectively. Visit THYQUIDITY.com for Full Prescribing Information.

To report SUSPECTED ADVERSE REACTIONS, contact Azurity Pharmaceuticals, Inc. at 1-800-461- 7449, or FDA at 1-800-FDA-1088 or www.fda.gov/MedWatch.

PP-THY-US-0006

 **THYQUIDITY™**
(levothyroxine sodium) oral solution, 100 mcg/5 mL

Azurity Pharmaceuticals, Inc. | 841 Woburn St., Wilmington, MA 01887 | Thyquidity.com
© 2022 Azurity Pharmaceuticals, Inc. All Rights Reserved. All Trademarks referred to are the property of their respective owners.
Product packaging and imagery are for representation purposes only and shall constitute the property of Azurity and all applicable affiliates.
PP-THY-US-0002