

## FOR YOUR PERSONAL DOSE, EVERY DAY

To manage your hypothyroidism appropriately, you need a specific amount of medicine daily.

THYQUIDITY is a liquid form of levothyroxine sodium, the most commonly prescribed medicine for hypothyroidism. THYQUIDITY allows your health-care provider to prescribe the dose that's just right for you. It is also designed to help you avoid complicated dosing schedules and pill splitting.

### FILL IN WITH YOUR HEALTHCARE PROVIDER...

As of today's date, \_\_\_\_/\_\_\_\_/\_\_\_\_, my daily THYQUIDITY dose is \_\_\_\_ ml

Note: If your prescriber changes your dose, cross out the old date and dose and enter the new date and dose.

#### DISCUSS WITH YOUR PHARMACIST...

Bring this sheet when picking up your THYQUIDITY prescription and discuss the following:



**Confirm that the pharmacist has provided a calibrated oral syringe based on your dose.**



**Ask your pharmacist to show you the line on the oral syringe where your liquid dose should reach.**



**Ask your pharmacist to review any foods, drinks, and medicines that may interfere with or should not be taken with THYQUIDITY.**

Note: THYQUIDITY comes in a package of two 100 mL bottles.

#### REVIEW AT HOME...

THYQUIDITY should be taken exactly as prescribed by your healthcare provider. Review and check off the following to help you stay on track.

- Take THYQUIDITY once a day (at the same time each day) on an empty stomach, 30 to 60 minutes before breakfast.
- Take THYQUIDITY at least 4 hours before or after taking medicine known to interfere with the absorption of THYQUIDITY, such as iron and calcium supplements and antacids.
- Confirm your prescribed dose before taking THYQUIDITY, and then locate the correct dose line on your oral syringe.
- Mark the date you open your THYQUIDITY bottle, and use within 8 weeks of opening.
- Take THYQUIDITY using only the oral syringe provided with your prescription by the pharmacy.
- Take THYQUIDITY by drawing the liquid into the oral syringe until it reaches your dose line exactly, and then administer directly into your mouth.
- Store THYQUIDITY at room temperature (68°F to 77°F) in the original bottle, and protect from light.
- Set a calendar reminder to refill your medicine a few days before you run out.

See if you are eligible to pay as little as \$0 for THYQUIDITY\*



Scan the code or visit  
[THYQUIDITYresources.com](https://www.thyquidity.com)

\*Offer valid for commercially insured patients only. Additional eligibility restrictions and terms and conditions apply.

#### SELECTED IMPORTANT SAFETY INFORMATION

- Thyroid hormones, including THYQUIDITY, should not be used for the treatment of obesity or for weight loss.
- Doses beyond the range of daily hormonal requirements may produce serious or even life-threatening manifestations of toxicity. (6, 10)

Please see additional Important Safety Information on the following page and full Prescribing Information including Boxed Warning at <https://www.thyquidity.com/pdf/Prescribing-Information.pdf>

**Hypothyroidism**

As a replacement therapy in primary (thyroidal), secondary (pituitary), and tertiary (hypothalamic) congenital or acquired hypothyroidism.

**Pituitary Thyrotropin (Thyroid Stimulating Hormone, TSH) Suppression**

As an adjunct to surgery and radioiodine therapy in the management of thyrotropin-dependent well-differentiated thyroid cancer.

Limitations of Use

- Not indicated for suppression of benign thyroid nodules and nontoxic diffuse goiter in iodine-sufficient patients.
- Not indicated for treatment of hypothyroidism during the recovery phase of subacute thyroiditis.

**IMPORTANT SAFETY INFORMATION****WARNING: NOT FOR TREATMENT OF OBESITY OR FOR WEIGHT LOSS**

*See full prescribing information for complete boxed warning*

- **Thyroid hormones, including THYQUIDITY, should not be used for the treatment of obesity or for weight loss.**
- **Doses beyond the range of daily hormonal requirements may produce serious or even life-threatening manifestations of toxicity. (6, 10).**

Do not use THYQUIDITY if you have uncorrected adrenal problems.

Thyroid hormones should be used with great caution in circumstances where the integrity of the cardiovascular system is suspected. In the elderly and in patients with cardiovascular disease, THYQUIDITY should be used with greater caution.

Tell your doctor about any other medical conditions you may have, especially heart disease, diabetes, blood clotting problems, and adrenal or pituitary gland problems. The dose of other drugs you may be taking to control these conditions may have to be changed while you are taking THYQUIDITY. If you have diabetes, check your blood sugar levels and/or the glucose in your urine, as ordered by your doctor, and immediately tell your doctor if there are any changes. If you are taking blood thinners, your blood clotting status should be checked often.

Taking too much levothyroxine has been associated with increased bone loss, especially in women after menopause.

Once your doctor has found your specific THYQUIDITY dose, it is important to have lab tests done, as ordered by your doctor, at least once a year.

Foods like soybean flour, cottonseed meal, walnuts, and dietary fiber may cause your body to absorb less THYQUIDITY from the gastrointestinal tract. Grapefruit juice may cause your body to absorb less levothyroxine and may reduce its effect. Let your doctor know if you eat these foods, as your dose of THYQUIDITY may need to be adjusted.

Use THYQUIDITY only as ordered by your doctor. Take THYQUIDITY as a single dose, preferably on an empty stomach, one-half to one hour before breakfast.

Products such as iron and calcium supplements and antacids can lower your body's ability to absorb levothyroxine, so THYQUIDITY should be taken 4 hours before or after taking these products.

Tell your doctor if you are pregnant or breastfeeding or are thinking of becoming pregnant while taking THYQUIDITY. Your dose of THYQUIDITY may need to be increased during your pregnancy.

Monitor your baby from birth to 3 months of age for vomiting and/or diarrhea as THYQUIDITY can cause gastrointestinal irritation due to the glycerol component.

It may take several weeks before you notice an improvement in your symptoms.

Tell your doctor about all medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements.

Tell your doctor or dentist that you are taking THYQUIDITY before any surgery.

Adverse reactions associated with THYQUIDITY therapy are primarily those of hyperthyroidism due to therapeutic overdosage: rapid or abnormal heartbeat, chest pain, difficulty catching your breath, leg cramps, headache, nervousness, irritability, sleeplessness, shaking, change in appetite, weight gain or loss, vomiting, diarrhea, increased sweating, difficulty tolerating heat, fever, changes in menstrual periods, swollen red bumps on the skin (hives) or skin rash, or any other unusual medical event. Tell your doctor if you develop any of these symptoms.

Partial hair loss may occur during the first few months you are taking THYQUIDITY.

***The Important Safety Information does not include all the information needed to use THYQUIDITY safely and effectively. Visit [THYQUIDITY.com](http://THYQUIDITY.com) for Full Prescribing Information.***

***To report SUSPECTED ADVERSE REACTIONS, contact Azurity Pharmaceuticals, Inc. at 1-800-461-7449, or FDA at 1-800-FDA-1088 or [www.fda.gov/MedWatch](http://www.fda.gov/MedWatch).***

PP-THY-US-0006

 **THYQUIDITY™**  
(levothyroxine sodium) oral solution, 100 mcg/5 mL

Azurity Pharmaceuticals, Inc. | 8 Cabot Road, Suite 2000, Woburn, MA 01801 | [Thyquidity.com](http://Thyquidity.com)  
© 2023 Azurity Pharmaceuticals, Inc. All Rights Reserved. All Trademarks referred to are the property of their respective owners. Product packaging and imagery are for representation purposes only and shall constitute the property of Azurity and all applicable affiliates.  
PP-THY-US-0007